

Cell Injury

By

Dr. Dina Abdallah

A central green rectangle is surrounded by six white ovals. The ovals are labeled: 'Mucin' (top), 'Gout' (top-left), 'Hyalinosis' (top-right), 'Amyloidosis' (bottom-right), 'Pathological calcification' (bottom), and 'Pathological pigmentation' (bottom-left). The 'Mucin' oval contains several small asterisk-like symbols.

Mucin

Gout

Hyalinosis

Intra and Extracellular
Accumulations and
Depositions

Pathological
pigmentation

Amyloidosis

Pathological
calcification

1-Deposition of Mucin

Normally mucin is secreted by mucous secreting cells e.g. Respiratory tract or GIT cells.

Excessive accumulation of mucopolysaccharides (muroid material) in:

- Epithelial cells is called "muroid degeneration"
- Connective tissue is called "myxomatous degeneration".

N/E:

Pale grey transparent slimy fluid.

M/E:

Pale blue with Hx.



1-Deposition of Mucin

A) Epithelial (muroid):



- E.g: Neoplastic cells: Mucin secreting tumors that occur in many organs e.g. ovary, colon, and stomach.



B) Connective tissue (myxomatous):

- E.g: Myxoma





Mucoid Adenocarcinoma





Myxoma



Mucin

Gout

Hyalinosis

Intra and Extracellular
Accumulations and
Depositions

Pathological
pigmentation

Pathological
calcification

Amyloidosis

2- Hyalinosis (Hyaline degeneration)

Def: Deposition of glassy, refractile homogenous structureless transparent material that stains red with eosin either inside cells or in connective tissue material.

Pathogenesis: Physico-chemical alteration of local cellular and extracellular protein with unclear mechanism.


Types:

- I) Intracellular
- II) Extracellular


Examples:

- 1- Corpora amylacea.
- 2- Russel bodies of plasma cells in Rhinoscleroma.
- 3- Mallory bodies in alcoholic hepatitis.


Corpora amylacia



Rhinoscleroma



Alcoholic hepatitis



Intra and Extracellular Accumulations and Depositions

```
graph TD; A([Mucin]) --- B[Intra and Extracellular Accumulations and Depositions]; B --- C([Hyalinosis]); C --- D([Amyloidosis]); D --- E([Pathological calcification]); E --- F([Pathological pigmentation]); F --- G([Gout]); G --- A;
```

Mucin

Hyalinosis

Amyloidosis

Pathological
calcification

Pathological
pigmentation

Gout

3-Gout (hyperuricaemia)

Def: Disturbance in purine metabolism with deposition of sodium urate in the tissue.

Causes:

a) Primary:

familial, ♂ > ♀ at 40 years due to increased purine breakdown or decreased clearance .

b) Secondary:

increased cellular destruction as in Polycythaemia rubra vera, leukaemia ... etc.

Ex: Joint:

- Recurrent acute attacks of arthritis: It affects metatarsophalangeal joint of big toe which become red, hot & swollen.



Mucin

Gout

Hyalinosis

Intra and Extracellular Accumulations and Depositions

Pathological
pigmentation

Pathological
calcification

Amyloidosis

AMYLOIDOSIS (AMYLOID DEGENERATION)

***Def.** Extra cellular deposition of abnormal proteinacious substance (amyloid protein) in many organs and tissues.

- **Morphological features:**

*** Grossly:** pale grayish brown, waxy and translucent material.

*** N/E:** (solid organs) as in liver, kidney & spleen show;


- **Shape** → preserved
- **Surface** → smooth
- **Size** → enlarged
- **Consistency** → firm & elastic
- **Color** → pale grayish brown & waxy.
- **C / S** → sharp edges, pale grayish brown & waxy.

*** M/E:**

- Hx&E-stained sections appears as homogenous Structure less pale red material (like fibrinoid necrosis).

**Thickened cortex with
waxy brown colour**





***Staining of amyloid material:**



I. Gross staining: staining of fresh slices of tissue.


- Lugol's iodine: → dark brown.
- Iodine + 1% sulphuric acid: → blue.

II. Microscopic staining: staining done on paraffin sections and frozen sections.

- Congo red stain: → orange red and apple green birefringence by polarized light.
- Metachromatic stains: methyl violet, gentian violet and crystal violet → rose red.







Crystal violet

Mucin

Gout

Hyalinosis

Intra and Extracellular
Accumulations and
Depositions

Pathological
pigmentation

Amyloidosis

Pathological
calcification

PATHOLOGICAL CALCIFICATION

***Def.** Pathological deposition of calcium salts (phosphate & carbonate) in the tissues (other than bone & teeth).

Types:

- **1. Dystrophic calcification**
-Occurs in nonviable tissue with normal blood calcium level
- **2. Metastatic calcification**
-Occurs in viable tissue with hypercalcemia
- **3. Stone formation**

***Types:**

I. Dystrophic calcification

***Def:** pathological deposition of Ca salts in degenerated or necrotic tissue with normal blood Ca level.

*** Sites:**

- Atheroma
- fibrosed valve
- old scar
- old thrombi,
- old infarct
- fat necrosis
- caseating T.B lesion
- pus of chronic abscess
- dead B ova or worm.



II. Metastatic calcification

*Def: pathological deposition of Ca salts in living tissues due to hypercalcaemia.





*Etiology: hypercalcaemia.

Causes:

- a) Excess absorption of calcium from the intestine as in;
 - Hypervitaminosis D (vitamin D intoxication)
 - Excess vitamin D and calcium in food of infants
- b) Excess mobilization of calcium from bone as:
 - Hyperparathyroidism & thyrotoxicosis
 - Prolonged immobilization in bed
 - Bone destruction by malignant tumors

*Sites:

- Walls of the lung alveoli.
- Mucosa of the stomach.
- Renal tubules.
- Arteries.



III. Stone formation.

- Ca salts deposition in the cavities of hollow organs
 - as - gall bladder
 - urinary tracts

Mucin

Gout

Hyalinosis

Intra and Extracellular Accumulations and Depositions

Pathological
pigmentation

Amyloidosis

Pathological
calcification

Pathological pigmentation

Pigments are colored substances that stain the tissue. These could accumulate intracellularly.

Types: may be exogenous or endogenous.

1- Exogenous Routes of entry:

- **Inhalation** as in pneumoconiosis due to silicosis & anthracosis.

- **Ingestion** as in chronic lead poisoning(plumbism).


- **Inoculation** as in tattooing

2- Endogenous

- **Melanin**

- **Lipofuscin (lipochrome)**

- **Hemosidrin**



Pathological pigmentation

2- Endogenous: Melanin

Melanin hyperpigmentation:

- 1- Exposure to sun: There is stimulation of MSH (melanocyte stimulating hormone).
- 2- Chronic irritation
- 3- Chloasma of pregnancy: Pigmentation of the face, nipple, and genitalia caused by ACTH, MSH and excess estrogen.
- 4- Addison's disease (chronic adrenal insufficiency) due to increase level of ACTH which has melanocytes stimulating effect.
- 5- Tumors:
 - Tumours of melanocytic origin;
 - a- Pigmented nevus. b- Malignant melanoma. .

Pathological pigmentation

2- Endogenous: Melanin

Melanin hypopigmentation:

- 1- Albinism: generalized hypopigmentation due to tyrosinase deficiency. Melanin pigment is absent from skin, hair and eyes.
 - 2- Leukoderma: congenital absence of melanocytes
 - 3- Vitiligo (acquired absence of melanocytes): either primary or secondary to skin lesions as: Scar, syphilis, and leprosy.
- Both leukoderma and vitiligo are patchy.

Albinism





Vitiligo

Pathological pigmentation

2- Endogenous: Lipochrome (Lipofuscin)


It is a yellowish brown fat soluble pigment. normally in the heart, testis, seminal vesicles, corpus luteum and adrenal cortex.


Causes:

1. Old age.
2. Atherosclerosis.
3. Wasting diseases.
4. Vitamin deficiency and severe malnutrition.
5. Cancer cachexia
6. Endocrinal disturbances

Pathological pigmentation

2- Endogenous: Lipochrome (Lipofuscin) Brown atrophy of the heart





Pathological pigmentation

2- Endogenous: Haemosiderosis

Hemosidrin is an iron containing pigment consisting of aggregates of ferritin

RBCs at the end of life span releases its HB content which is then metabolized into :


1- Globulin (that is reused).

2- Haem which consists of:

- Fe (that is reused)

- Biliverdin → bile.

- -Fe is absorbed from duodenum & carried in plasma transferritin to be stored as Fe or apoferritin in macrophages in liver , spleen & BM.



A

